



# Analysis on the relevance between periodontal inflammation and macrophage polarization and function in gingival tissue



Lina Zhou<sup>1</sup>, Honglei Qu<sup>1</sup>, Xuan Li<sup>1</sup>, Lijuan Sun<sup>1</sup>, Faming Chen<sup>1\*</sup>,

<sup>1</sup>State Key Laboratory of Military Stomatology, National Clinical Research Center for Oral Diseases, Shaanxi Engineering Research Center for Dental Materials and Advanced Manufacture, Department of Periodontology, Stomatology hospital, Fourth Military Medical University, Xi'an 710032, China

**Background:** Activated macrophages play a central role in the development progress of chronic periodontitis. Polarized macrophages M1- and M2-type macrophages have distinct effects on periodontal inflammation through secreting various cytokines. Our study aims to observe the M1/M2 ratios and expression of macrophage-related cytokines in different periodontal status, then investigate the relevance between periodontal inflammation and macrophage polarization and function.

**Methods:** Forty-three gingival samples were collected from patients with chronic periodontitis (n=13), gingivitis (n=19) and periodontal health individual (n=11). Periodontal index probing depth (PD), clinical attachment level (CAL) and gingival index (GI) were recorded before collection. M1/M2 ratios were detected through immunofluorescence staining, while the expression levels of macrophage-related cytokines TFN- $\alpha$ , IFN- $\gamma$ , IL-6, IL-12, IL-4 and IL-10 were observed through immunohistochemistry staining. The relevance between them was analysed by statistical software.

**Results:** Compared with periodontal health group, chronic periodontitis group has a significant increase in M1/M2 ratio accompanied by enhanced TFN- $\alpha$ , IFN- $\gamma$ , IL-6 and IL-12 expression (p<0.05), while gingivitis group only has enhancements in TNF- $\alpha$ , IL-12 and IL-4 expression (p<0.05), without any difference of M1/M2 ratio. Both M1/M2 ratio, IFN- $\gamma$  and IL-6 expression levels in chronic periodontitis group were significant higher than gingivitis group (p<0.05). Meanwhile, M1/M2 ratios were positively correlated with PD, IFN- $\gamma$  and IL-6 (p<0.01), PD was positively correlated with IL-6 and IFN- $\gamma$  and negatively correlated with IL-4 (p<0.05).

**Conclusion:** Transformation of M1 and M2 has a crucial impact on periodontal inflammation. Predominance of M1-type macrophages and M1-related cytokines associates with the pathogenesis of chronic periodontitis. Regulation of macrophages polarization may be a potential therapeutic target for immunomodulatory therapy of chronic periodontitis.

## Results




Fig.1 Samples collection and HE staining of gingival tissues




Fig. 2 Expression of M1- and M2-type macrophages in gingival tissues




Fig. 3 Expression of cytokines IFN- $\gamma$  and TNF- $\alpha$  in gingival tissues




Fig. 4 Expression of cytokines IL-6 and IL-12 in gingival tissues




Fig. 5 Expression of cytokines IL-4 and IL-10 in gingival tissues




Fig. 6 Correlation between periodontal index and M1/M2 ratios and macrophage-related cytokines